

VDI References

Cited Chronologically

Hyperlinked references– click on reference to review PubMed abstract

Bolded references are more noteworthy

Thymidine Kinase

1. Gronowitz JS, Hagberg H, Källander CFR, Simonsson B (April 1983). "The use of serum deoxythymidine kinase as a prognostic marker, and in the monitoring of patients with non-Hodgkin's lymphoma". *Br. J. Cancer* 47 (4): 487–95. doi:10.1038/bjc.1983.78.PMC 2011337. PMID 6849793.
2. Hagberg H, Glimelius B, Gronowitz JS, Killander A, Källander CFR, Schröder T (July 1984). "Biochemical markers in non-Hodgkin's lymphoma stages III and IV and prognosis: a multivariate analysis". *Scand J Haematol* 33 (1): 59–67.
3. **Greenlee PG, et al. Lymphomas in dogs. A morphologic, immunologic, and clinical study. *Cancer* (1990). 66.3: 480–490.**
4. Gronowitz, J S. et al. 1996. "Thymidine Kinase: Biochemical Background and Clinical Applications: Background and use in gene therapy." Biovica AB. Uppsala, Sweden.
5. Gronowitz, J S. et al. 1996. "Thymidine Kinase: Biochemical Background and Clinical Applications: The quantitative assay for TK and its clinical use." Biovica AB. Uppsala, Sweden.
6. Gronowitz, J S. et al. 1996. "Thymidine Kinase: Biochemical Background and Clinical Applications: Use of TK assay in serum, plasma and other body fluids." Biovica AB. Uppsala, Sweden.
7. **Nakamura, et al, "Plasma Thymidine Kinase Activity in Dogs with Lymphoma and Leukemia", 1997. *J Vet Med Sci.* 59(10): 957-960**
8. Tams, Todd R. "Inflammatory Bowel Disease and Intestinal Lymphoma in Cats." Presented at Atlantic Coast Veterinary Conference. (2001).
<http://www.vin.com/vindbpub/searchpb/proceedings/pr05000/pr00427.htm>
9. Little, S (2002) "Feline Inflammatory Bowel Disease." The Winn Feline Foundation.
http://www.winnfelinehealth.org/Pages/Feline_IBD_Web.pdf.
10. **Von Euler, et al, "Serum Thymidine Kinase Activity in Dogs with Malignant Lymphoma: A Potent Marker for Prognosis and Monitoring the Disease", 2004. *J Vet Intern Med.* 18: 696-702**
11. **Madewell, B. R. (2004), Editorial: Serum Thymidine Kinase Activity: An Alternative to Histologic Markers of Cellular Proliferation in Canine Lymphoma. *Journal of Veterinary Internal Medicine,* 18: 595–596.**

VDI References

12. Von Euler, et al, "A Non-Radiometric Method for Measuring Serum Thymidine Kinase Activity in Malignant Lymphoma in Dogs", 2006. *Res Vet Sci.* **80: 17-24**
13. Evans, S E. et al. "Comparison of endoscopic and full-thickness biopsy specimens for diagnosis of inflammatory bowel disease and alimentary tract lymphoma in cats." *JAVMA.* (2006) 229.9: 1447-1450.
14. Lana S E., Rutteman G R., Withrow S J. (2007). *Canine lymphomas In: Small animal clinical oncology: Withrow S J., Vail D M. 4th edition, Elsevier Saunders 699-728pp.*
15. Von Euler, et al, "Monitoring Therapy in Canine Malignant Lymphoma and Leukemia with Serum Thymidine Kinase 1 Activity - Evaluation of a New, Fully Automated Non-Radiometric Assay", 2008. *Int J Onc.* **34:505-510**
16. Argyle, D J. et al. "Decision Making in Small Animal Oncology". (2008) Wiley-Blackwell. 197-225pp.
17. Wilson, Heather M. "Feline Alimentary Lymphoma: Demystifying the Enigma." *Topics in Companion Animal Medicine.* (2008) Vol 23(4): 177-184.
18. Selting, et al, "Serum Thymidine Kinase Concentrations in Hemangiosarcoma versus Healthy Dogs", 2009. Abstract presented at **Veterinary Cancer Society 2009 Annual Conference**
19. Hall, E J. (2009) "Inflammatory bowel disease in dogs and cats." Hills's Pet Nutrition. <https://protrain.hs.llnwd.net/e1/.../GI%20technical%20booklet.pdf>.
20. Sakamoto, L. et al. "Serum Thymidine Kinase Activity as a Useful Marker for Bovine Leukosis." *J Vet Diagn Invest.* (2009). 21(6): 871-4.
21. Henry C J, Higginbotham M L. (2010). *Cancer Management in Small Animal Practice.* Elsevier Saunders. 1-14, 28-34, 41-100, 136-145, 249-264, 343-355pp.
22. Washabau, R., et al. The WSAVA International Gastrointestinal Standardization Group. (2010). Endoscopic, Biopsy, and Histopathologic Guidelines for the Evaluation of Gastrointestinal Inflammation in Companion Animals. *Journal of Veterinary Internal Medicine*, 24: 10–26. doi: 10.1111/j.1939-1676.2009.0443.x
23. Zwingenberger, A. L. et al. "Ultrasonographic Evaluation of the Muscularis Propria in Cats with Diffuse Small Intestinal Lymphoma or Inflammatory Bowel Disease." *J Vet Intern Med* (2010) 24: 289-292.
24. Chen, Y. et al. "Serum Thymidine kinase 1 correlates to clinical stages and clinical reactions and monitors the outcome of therapy of 1,247 cancer patients in routine clinical settings." *Int J Clin Oncol* (2010). 15:359-368.
25. Pan, Zhu-Lin. Et al. "Serum Thymidine kinase 1 concentration as a prognostic factor of chemotherapy-treated non-Hodgkin's lymphoma patients." *J Cancer Res Clin Oncol.* (2010). DOI 10.1007/s00432-010-0769-z.

VDI References

26. He E, Xu XH, et al. Thymidine kinase1 is a potential marker for prognosis and monitoring the response to treatment of patients with breast, lung and esophagus cancer and non-Hodgkin's lymphoma. *Nucleoside, Nucleotides, Nucleic Acids* (2010); 29: 352–359.
27. Thamm,D., et al. 2011. Serum thymidine kinase activity in canine splenic hemangiosarcoma. Abstract presented at the American College for Veterinary Internal Medicine 2011 annual conference.
28. Scott, K., Zoran, D., Mansell, J., Norby, B. and Willard, M. (2011), Utility of Endoscopic Biopsies of the Duodenum and Ileum for Diagnosis of Inflammatory Bowel Disease and Small Cell Lymphoma in Cats. *Journal of Veterinary Internal Medicine*, 25: 1253–1257. doi: 10.1111/j.1939-1676.2011.00831.x
29. Kiupel, M. et al. “Diagnostic Algorithm to Differentiate Lymphoma From Inflammation in Feline Small Intestinal Biopsy Samples.” *Vet Pathol.* (2011). 48: 212.
30. Barr F., Gaschen L. “BSAVA Manual of Canine and Feline Ultrasonography.” BSAVA 2011. England. 100-109, 124-146pp
31. **Thamm, et al, (2011), Elevated serum thymidine kinase activity in canine splenic hemangiosarcoma. *Veterinary and Comparative Oncology*. doi: 10.1111/j.1476-5829.2011.00298.x**
32. **Taylor, S., et al. Serum thymidine kinase activity in clinically healthy and diseased cats: a potential biomarker for lymphoma. *Journal of Feline Medicine and Surgery*. doi: 10.1177/1098612X12463928**
33. **Selting,K., et al. 2012. Thymidine kinase 1 and c-reactive protein as biomarkers for screening clinically healthy dogs for occult disease. Abstract/ oral presentation presented at the American College for Veterinary Internal Medicine 2012 annual conference.**
34. **VDI Laboratory, 2012. The Application of Serum Thymidine Kinase Type 1 Concentration as a Prognostic Indicator for Canine Lymphoma and Hemangiosarcoma. Poster presented at the Veterinary Cancer Society 2012 annual conference.**
35. **VDI Laboratory, 2012. Evaluation of Serum Thymidine Kinase Type 1 as a Diagnostic Biomarker in the Differentiation of Feline Lymphoma from Inflammatory Bowel Disease. Poster presented at the Veterinary Cancer Society 2012 annual conference.**
36. **VDI Laboratory. 2012. Case Studies of the Utility of VDI-TKcanine+ and VDI-TKfeline for Cancer Diagnosis and Therapeutic Monitoring. Poster presented at the Veterinary Cancer Society 2012 annual conference.**
37. **Westberg,S., et al. 2012. Correlation of pretreatment Thymidine Kinase concentration with remission duration and survival in dogs with lymphoma. Presented at the World Veterinary Cancer Congress 2012 annual conference.**
38. **Selting,K., et al. 2013. Serum thymidine kinase 1 and C-reactive protein as biomarkers for screening clinically healthy dogs for occult disease. *Veterinary and Comparative Oncology*. doi: 10.1111/vco.12052**

VDI References

39. Selting,K., et al. 2013 Use of thymidine kinase type 1 and c-reactive protein to detect cancer in dogs. Abstract/ oral presentation presented at the Veterinary Cancer Society 2013 annual conference.

Acute Phase Proteins (C-Reactive Protein, Haptoglobin)

40. Dillman,R.C. and E.H.Coles, 1966. A canine serum fraction analogous to human C-reactive protein. Am.J.Vet.Res. 27:1769-75.
41. Riley,R.F. and Coleman M.K.. Isolation of C-reactive proteins of man monkey, rabbit and dog by affinity chromatography on phosphorylated cellulose. Clin.Chim.Acta. 1970;30:483-96.
42. Riley,R.F. and W.Zontine, 1972. Further observations on the properties of dog C-reactive protein and the C-reactive protein response in the dog. J.Lab.Clin.Med. 80:698-703.
43. Kushner,I. and G.Feldmann, 1978. Control of the acute phase response. Demonstration of C-reactive protein synthesis and secretion by hepatocytes during acute inflammation in the rabbit. J.Exp. Med. 148: 466-77.
44. Hulton,N.R., D.J.Johnson and D.W.Wilmore, 1985: Limited effects of prostaglandin inhibitors in Escherichia coli sepsis. Surgery 98:291-7.
45. Caspi,D., F.W.J.J.Snel, et al. 1987. C-reactive protein in dogs. Am.J.Vet.Res. 48:919-21.
46. Eckersall,P.D., M.J.A.Harvey, et al. 1993. Acute phase proteins in canine pregnancy (canis familiaris). J.Reprod.Fert. S47:159-64.
- 47. Yamamoto,S., T.Shida, et al. 1993. Changes in serum C-reactive protein levels in dogs with various disorders and surgical traumas. Vet.Res.Com.17:85-93.**
48. Yamamoto,S., T.Shida, et al. 1994. Determination of C-reactive protein in serum and plasma from healthy dogs and dogs with pneumonia by ELISA and slide reversed passive latex agglutination test. Vet. Quart. 16:74-7.
49. Yamamoto,S., T.Shida, et al. 1994. Serum C-reactive protein and immune responses in dogs inoculated with Bordetella bronchiseptica (phase I cells). Vet.Res.Com. 18:347-57.
50. Otabe,K., T.Ito, et al, 2000. C-reactive protein (CRP) measurement in canine serum following experimentally induced acute gastric mucosal injury. Lab.Anim. 34:434-8.
- 51. Hayashi,S., T.Jinbo, et al. 2001. A comparison of the concentrations of C-reactive protein and alpha1-acid glycoprotein in the serum of young and adult dogs with acute inflammation. Vet.Res.Com. 25:117-26.**
52. Shimada,T., Y.Ishida, et al. 2002. Monitoring C-reactive protein in beagle dogs experimentally inoculated with Ehrlichia canis. Vet.Res.Com 26:171-7.

VDI References

53. **Martínez-Subiela,S., F.Tecles, P.D.Eckersall and J.J.Ceron, 2002: Serum concentrations of acute phase proteins in dogs with leishmaniasis. Vet. Rec. 150:241-4.**
54. Spillmann,T., J.Korrell, et al. 2002. Serum canine pancreatic elastase and canine C-reactive protein for the diagnosis and prognosis of acute pancreatitis in dogs. J.Vet.Int.Med. 16:635.
55. Kuribayashi,T., T.Shimada, et al. 2003. Determination of serum C-reactive protein (CRP) in healthy beagle dogs of various ages and pregnant beagle dogs. Exp. Animals 52: 387-90.
56. Borer,L.R., J.E.Peel, et al., 2003. Effect of carprofen, etodolac, meloxicam, or butorphanol in dogs with induced acute synovitis. Am.J.Vet.Res. 64:1429-37.
57. Martínez-Subiela,S., P.J.Ginel and J.J.Ceron, 2004. Effects of different glucocorticoid treatments on serum acute phase proteins in dogs. Vet.Rec. 154: 814-7.
58. **Kjelgaard-Hansen, M. 2004. Canine C-reactive protein – a study on the applicability of canine serum C-reactive protein. PhD thesis, The Royal Veterinary and Agricultural University, Frederiksberg,Denmark.**
59. K. Kriz, F. Ibraimi, M. Lu, et al. 2005. Detection of C-reactive Protein utilizing magnetic permeability detection based immunoassays. Anal.Chem.77(2005) 5920.
60. **Kjelgaard-Hansen M, Jensen AL et al. 2006. Use of serum C-reactive protein as an early marker of inflammatory activity in canine type II immune-mediated polyarthritis: case report. Acta Vet Scand.48:9.**
61. **Matijatko, V., V. Mrljak et al. 2007. Evidence of an acute phase response in dogs naturally infected with Babesia canis. Vet. Parasitol. 144:242-50.**
62. **Mansfield, C.S., F.E. James and I.D. Robertson. 2008. Development of a clinical severity index for dogs with acute pancreatitis. J. Am.Vet.Med.Assoc. 233:936-44.**
63. **Nakamura, M., M.Takahashi, et al. 2008. C-reactive protein concentration in dogs with various diseases. J.Vet.Med.Sci. 70:127-31.**
64. K. Kriz, L-O. Hansson, D. Kriz, 2008. CRP: an important biomarker for infectious disease and its role in POC. CLI. Apr 2008.
65. **Dabrowski R., K. Kostro et al. 2009. Usefulness of C-reactive protein, serum amyloid A component, and haptoglobin determinations in bitches with pyometra for monitoring early post-ovariohysterectomy complications.Theriogenology 72:471-6.**
66. **Griebsch C., G. Arndt et al. 2009. C-reactive protein concentration in dogs with primary immune-mediated hemolytic anemia. Vet.Clin.Pathol. 38:421-5.**
67. Yu, G., Y. Yu et al. 2009. Effect of periodontitis on susceptibility to atrial fibrillation in an animal model. J. Electrocardiol. 43:359-66.

VDI References

68. **Lowrie, M., J. Penderis, et al 2009. The role of acute phase proteins in diagnosis and management of steroid-responsive meningitis arteritis in dogs. Vet. J. 182:125-30.**
69. Ibraimi F., Kriz K., Merin H., Kriz, D. et al. 2009. Magnetic permeability based diagnostic test for the determination of the canine C-reactive protein concentration in undiluted whole blood. JMMM. 321(2009) 1632-1634
70. Tvarijonaviciute A., S. Martinez et al. 2010. Serum acute phase protein concentrations in dogs during experimentally short-term induced overweight. A preliminary study. Res.Vet.Sci. Epub ahead of print PMID: 20553703
71. Wakshlag JJ, Stokol T et al. 2010. Evaluation of exercise-induced changes in concentrations of C-reactive protein and serum biochemical values in sled dogs completing a long-distance endurance race. Am.J.Vet.Res. 71:1207-13
72. **Kocaturk M., S. Martinez, et al. 2010. Prognostic value of serum acute-phase proteins in dogs with parvoviral enteritis. J. Small. Anim. Pract. Epub ahead of print PMID: 20630018**
73. **Jergens, A.E., J. Crandell, et al. 2010. Comparison of oral prednisone and prednisone combined with metronidazole for induction therapy of canine inflammatory bowel disease: a randomized controlled trial. J. Vet. Intern. Med. 24:269-77.**
74. Yuki, M. and T. Hirano. 2010. Use of a combination of prednisolone and rosuvastatin for treatment of a pyogranuloma in a dog. J Am Vet Med Assoc. 236:767-9.